

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

382 REVIEWS

Geological Report on Arenac County. By W. M. GREGORY. Michigan Geol. and Biol. Surv. Publ. 11, Geol. Series 8. 1911. Pp. 148, pls. 6, figs. 18, map 1.

The county is underlain by Mississippian and Pennsylvanian formations which dip slightly to the south. The economic resources are very slight. Limestone, gypsum, and clay are of local usefulness.

T. T. Q.

Annotated Bibliography of Iowa Geology and Mining. By Charles Keyes. Iowa Geol. Surv., Vol. XXII. 1913. Pp. 908.

Most of the first 150 pages of the bibliography are given to historical material. The bibliography is arranged alphabetically under the names of authors and subjects.

T. T. Q.

Prince George's County. By William Bullock Clark. Maryland Geol. Surv. 1911. Pp. 251, pls. 13, figs. 3. Accompanied by Prince George's County Atlas, 2 maps.

The sixth of a series of reports dealing with the physical features of the several counties of Maryland. A full discussion of the stratigraphic geology of the county accompanies a description of the physiography, mineral resources, soils, forests, climate, and hydrography.

The geologic formations represented in the county range from Archean to very recent. After the Archean, no formations are represented below the Potomac group of the Comanchean. Later formations represent the Cretaceous, the Eocene (Pamunkey), the Miocene (Chesapeake), and later periods.

T. T. Q.

The Manhattan Schist of Southeastern New York State and Its Associated Igneous Rocks. By Charles Reinhard Fettke. A dissertation (Columbia). Annals N.Y. Acad. Sci., XXIII, April 30, 1914, pp. 193–260, Plates VIII–XV.

The erosion of northeast-southwest trending anticlines and synclines has exposed the Manhattan schist in a series of roughly parallel belts south of the Highlands of the Hudson and east of the Hudson River. The Manhattan is a quartz-mica-feldspar schist and the young-